

The dangers of extremes

Donald Marolf

*Physics Department, UCSB, Santa Barbara,
CA 93106, USA
marolf@physics.ucsb.edu*

Essay written for the Gravity Research Foundation 2010 Awards for Essays on
Gravitation

Submission date: March 28, 2010

ABSTRACT: While extreme black hole spacetimes with smooth horizons are known at the level of mathematics, we argue that the horizons of physical extreme black holes are effectively singular. Test particles encounter a singularity the moment they cross the horizon, and only objects with significant back-reaction can fall across a smooth (now non-extreme) horizon. As a result, classical interior solutions for extreme black holes are theoretical fictions that need not be reproduced by any quantum mechanical model. This observation suggests that significant quantum effects might be visible outside extreme or nearly extreme black holes. It also suggests that the microphysics of such black holes may be very different from that of their Schwarzschild cousins.

Contents

1. Introduction and review	1
2. Falling into extreme black holes	3

1. Introduction and review

For reasons related to the strict vanishing of their temperature, extreme black holes are generally considered impossible to form in classical physics. While any non-extremality parameter can be made small with enough care, it cannot be brought to zero in finite time. Yet precisely extreme black holes can in principle be formed through quantum processes. For example, extreme Reissner-Nordström (RN) black holes can be pair created from the vacuum by strong electric and magnetic fields [1]. Extreme magnetically-charged such black holes may even be stable in our universe. More generally, in theories of gravity with appropriate supersymmetries, Hawking radiation can cause charged black holes to decay to extremality in finite time. In this sense precisely extreme black holes can be said to be “physical,” at least in certain theories¹.

In the supersymmetric context, extreme solutions have provided a marvelous laboratory in which to explore fundamental issues. In particular, within string theory a detailed accounting of the Bekenstein-Hawking entropy has been given in terms of D-brane configurations for many extreme and nearly extreme black holes following [2]. We also mention the research program associated with Mathur [3] which attempts to replace black holes by collections of horizon-free geometries, and which has also focused on extremal settings.

Such works emphasize extreme black holes for which the maximal analytic extension of the stationary exterior has a smooth horizon where curvatures are small. Typical examples resemble extreme Reissner-Nordström, whose conformal diagram is shown in figure 1. It may thus appear natural to ask whether these microscopic models also describe the region marked B inside the horizon. This can be a large region in which curvatures are small so long as one stays well away from the singularity.




Figure 1: The conformal diagram for an extreme RN black hole. Future and past null infinity ($I^\pm$) are shown for two asymptotic regions, A and C. The dotted line is a time-like singularity and the dashed lines are Cauchy horizons. The two circles are internal infinities lying at the ends of infinite throats.

¹Since we know of no mechanism to stabilize extreme Kerr against quantum mechanical radiation of gravitons, this essay will focus on spherically-symmetric extreme black holes. We also take “black holes” to have compactly-generated horizons.

Nevertheless, we argue below that physical extreme black holes simply have no analogue of region B. Instead, although curvatures are small everywhere outside the horizon, test particles encounter a curvature singularity the instant they enter the black hole. The physical extreme spacetime is better described by figure 2, in which the horizon has been replaced by a null singularity. Infalling objects that produce large back-reaction effects may experience a smooth horizon, but one which is now non-extreme and the region they enter has properties quite different from region B.

Our argument relies on the mass-inflation instability of the inner horizon for non-extreme RN black holes, which we now review. The conformal diagram for the analytic extension of static non-extreme RN is shown in figure 3. Recall that region A_3 is separated from regions B_1, B_2 by Cauchy horizons, which are also the inner horizons of the black hole. One sees from the diagram that initial data in A_1, A_2 does indeed determine evolution in region A_3 , but not in B_1, B_2 .




Figure 2: The effective conformal diagram for physical extreme RN black holes. The dotted line is a null singularity.




Figure 3: The analytic extension of a non-extreme RN black hole. Again, the dotted vertical lines are timelike singularities and the dashed lines are Cauchy horizons. The fine diagonal dash-dotted lines represent a series of constant-phase surfaces for a radial ingoing spherical wave. The wave undergoes an infinite number of oscillations below I^+ , and thus below the Cauchy horizon. But the proper distance (or affine parameter) across the wavefronts to the Cauchy horizon is finite. Thus, any observer falling through the Cauchy horizon would experience an infinite number of oscillations in a finite time; i.e., the radiation has been infinitely blueshifted

Recall also the infinite blueshift between I^+ and the Cauchy horizon, as illustrated in figure 3. It was suggested long ago by Penrose [4] that this blueshift could make the Cauchy horizon unstable. Tiny bits of (e.g.) gravitational radiation produced by an infalling object would lead to a divergence, replacing the Cauchy horizon with a singularity. This scenario was confirmed by Poisson and Israel [5] and rigorously demonstrated by Dafermos [6], though the presence of some right-moving perturbation also turns out to be important.

The full picture [7] appears to be that shown in figure 4. Any perturbation triggers an instability which forms a singularity. One piece of the singularity is null and simply replaces the would-be Cauchy horizon. Along this null singularity the area of the (singular) null congruence shrinks to zero, where it joins a spacelike singularity. Similar results appear to hold for rotating black holes [8], though the non-spherical setting is more difficult to investigate.

2. Falling into extreme black holes

It is interesting to ask how far the new singularity of figure 4 lies inside the horizon, especially in the extreme limit where the inner horizon (at area-radius r_-) of the unperturbed hole coincides with the outer horizon (at area-radius r_+). For simplicity, we concentrate on black holes with two asymptotic regions as shown in figures 3 and 4. Black holes that form by collapse should behave similarly, at least as probed by observers who fall into the black hole long after it has formed.

To give meaning to the above question, consider a freely falling observer in the unperturbed spacetime of figure 3 who enters from region A_1 with energy per unit mass $\gamma = -U \cdot \partial_t$, where U^a is the observer's 4-velocity and ∂_t is the usual static Killing field of the exterior. This observer requires a proper time $\Delta\tau$ to fall from r_+ to r_- where

$$\Delta\tau = \frac{r_+ - r_-}{\gamma} + \mathcal{O}((r_+ - r_-)^3/r_+^2\gamma^3) \quad (2.1)$$

with $c = 1$.

Now consider a freely-falling observer satisfying the same initial conditions in the perturbed RN spacetime of figure 4. No matter how small the perturbation at r_+ , before the proper time (2.1) elapses for our observer this perturbation must grow large, else the original cauchy horizon would remain. But once the perturbation is large, the spacetime quickly evolves toward a singularity. Our infalling observer must therefore reach the singularity in a time of order (2.1) after crossing the horizon. Again, this conclusion holds *no*




Figure 4: A perturbed non-extreme RN black hole. The perturbation triggers an instability which generates the singularity indicated by the dotted line. The singularity may have both null and spacelike parts, though for a black hole with two asymptotic regions there need be no spacelike part if the perturbation is sufficiently weak. No Cauchy horizons remain.

matter how small the initial perturbation, though in practice there will of necessity be perturbations of some minimal size set by quantum fluctuations.

For observers with fixed γ , the time (2.1) vanishes in the extreme limit $r_- \rightarrow r_+$. Their experiences in this limit are thus described by figure 2, in which the singularity coincides with the outer horizon. In this sense, figure 2 describes the extreme limit of a physical RN black hole².

Suppose now that one is given a precisely extreme black hole, created perhaps through one of the quantum processes mentioned in the introduction. Any observer falling into the black hole will carry with them a small amount of energy. Because the horizon is compactly-generated, any such perturbation will make the black hole non-extreme. It follows that the experience of any observer (with small but non-zero back-reaction) falling into an extreme black hole is given by the extreme limit of their experience falling into non-extreme black holes³.

The details of this experience are determined by the observer's back-reaction. The larger the resulting non-extremality, the more proper time may pass before encountering a singularity. However, defining the test particle limit as the limit of zero back reaction, one may say that test particles experience a singularity the moment they cross the original horizon.

As noted above, the situation may be somewhat different if the extreme black hole forms from decay of a single-asymptotic-region black hole, but only as probed by observers who fall in soon after the black hole has formed. Late time observers falling into the black hole must again quickly encounter a singularity. We expect this situation to be described by figure 5, which as described in the caption approximates figure 2 from the view-

point of late-time observers. In particular, the classical interior solution described by region B of figure 1 is a theoretical fiction that does not describe the experience of any observer.

It is interesting to ponder the implications for quantum discussions of black holes. First and foremost, the presence of strong curvatures (and strong curvature gradients) *immediately* inside the horizon suggests that significant quantum effects might possibly be visible outside extreme black holes, perhaps in scattering processes involving very low energy quanta as these events would be associated with minimal back-reaction. While we have nothing specific to propose at this time, there may be a window of opportunity, especially for non-perturbative effects. It is also conceivable that such non-perturbative

²Strictly speaking, the internal infinity of figure 2 is removed by whatever process creates the extreme black hole; e.g., decay from non-extremality by Hawking radiation. One might say that the infinite throat has been capped. However, this cap recedes with time so that an internal infinity in some sense emerges after a long time has passed.

³There is no corresponding argument for non-compact extreme horizons such as the Poincaré horizon of anti-de Sitter space. In such cases, non-extreme versions of the original horizon may not even satisfy the same boundary conditions and there is no reason to expect continuity.




Figure 5: The conjectured diagram for extreme black holes formed by decay of non-extreme black holes with a single asymptotic region. The conformal structure resembles that of Schwarzschild. However, the time-translation symmetry of the exterior will be strongly broken in the interior (shaded) so that the time required for $\gamma = 1$ observers to fall from the horizon to the singularity will vanish in the late-time limit. Observers who fall in long after formation will effectively experience fig. 2.

quantum effects could provide large corrections to the entropy of extreme black holes⁴.

In addition, the above observation alters our expectations for microscopic descriptions of extreme black holes. This is good news for many proposed models. It is not hard to imagine that an object's collision with either a stack of D-branes or with the Planck-curvature fuzz suggested by Mathur's program [3] would destroy the object in a manner that might resemble the collision with a singularity. However, since the horizons of extreme black holes are so different from those of their Schwarzschild cousins, this victory is somewhat hollow. The challenge for such models is merely postponed to the difficult regime of highly excited states, with large energies above extremality, where the models must somehow reproduce the gentle experience of an observer crossing a large smooth horizon. Since the macrophysics is very different in this regime, appropriate microscopic models may be very different as well.

Acknowledgements

It is a pleasure to thank Patrick Brady, Gary Horowitz, Harvey Reall, and Eric Poisson for numerous discussions over many years of both extreme and non-extreme black holes. This work was supported in part by the National Science Foundation under Grant No PHY08-55415, and by funds from the University of California.

References

- [1] F. Dowker, J. P. Gauntlett, S. B. Giddings and G. T. Horowitz, "On pair creation of extremal black holes and Kaluza-Klein monopoles," *Phys. Rev. D* **50**, 2662 (1994) [arXiv:hep-th/9312172].
- [2] A. Strominger and C. Vafa, "Microscopic Origin of the Bekenstein-Hawking Entropy," *Phys. Lett. B* **379**, 99 (1996) [arXiv:hep-th/9601029].
- [3] S. D. Mathur, "The fuzzball proposal for black holes: An elementary review," *Fortsch. Phys.* **53**, 793 (2005) [arXiv:hep-th/0502050].
- [4] R. Penrose, in *Battelle Rencontres*, edited by C. M. DeWitt and J. A. Wheeler (W.A. Benjamin, New York, 1968) pg 222.
- [5] E. Poisson and W. Israel, "Internal structure of black holes," *Phys. Rev. D* **41**, 1796 (1990).
- [6] M. Dafermos, "The interior of charged black holes and the problem of uniqueness in general relativity," *Communications on Pure and Applied Mathematics*, **58**, 445-504, (2005) [arXiv:gr-qc/0307013].
- [7] P. R. Brady and J. D. Smith, "Black Hole Singularities: A Numerical Approach," *Phys. Rev. Lett.* **75**, 1256 (1995) [arXiv:gr-qc/9506067].
- [8] A. Ori, "Structure Of The Singularity Inside A Realistic Rotating Black Hole," *Phys. Rev. Lett.* **68**, 2117 (1992); P. R. Brady, S. Droz and S. M. Morsink, "The late-time singularity inside non-spherical black holes," *Phys. Rev. D* **58**, 084034 (1998) [arXiv:gr-qc/9805008].

⁴Since the spacetime outside is not significantly disturbed, one would not expect large perturbative quantum effects. Indeed, perturbative analyses of quantum corrections to extreme black holes indicate no such effects [9].

- [9] D. A. Lowe, “Comment on ‘Do Semiclassical Zero Temperature Black Holes Exist’,” Phys. Rev. Lett. **87**, 029001 (2001) [arXiv:gr-qc/0011053].